

ON MODULAR BALL-QUOTIENT SURFACES OF KODAIRA DIMENSION ONE

ALEKSANDER MOMOT

ABSTRACT. Let $\Gamma \subset \mathbf{PU}(2, 1)$ be a lattice which is not co-compact, of finite Bergman-covolume and acting freely on the open unit ball $\mathbf{B} \subset \mathbb{C}^2$. Then the compactification $X = \overline{\Gamma \backslash \mathbf{B}}$ is a projective smooth surface with an elliptic compactification divisor $D = X \setminus (\Gamma \backslash \mathbf{B})$. In this short note we discover a new class of unramified ball-quotients X . We consider ball-quotients X with $kod(X) = h^1(X, \mathcal{O}_X) = 1$. We prove that each minimal surfaces with finite Mordell-Weil group in the class described is up to an étale base change the pull-back of the elliptic modular surface which parametrizes triples (E, x, y) of elliptic curves E with 6-torsion points $x, y \in E[6]$ such that $\mathbb{Z}x + \mathbb{Z}y = E[6]$.

1. INTRODUCTION

Let the symbol \mathcal{T} denote the class of complex projective smooth surfaces X which contain pairwise disjoint elliptic curves D_1, \dots, D_{h_X} such that $U = X \setminus \bigcup D_i$ admits the open unit ball $\mathbf{B} \subset \mathbb{C}^2$ as universal holomorphic covering; as explained in [7], \mathcal{T} forms the 'generic' class of compactified ball-quotient surfaces. There are several motivations to study surfaces in \mathcal{T} without assuming that $\pi_1(U, *)$ with its Poincaré action on \mathbf{B} is an arithmetic lattice of $\mathbf{PU}(2, 1)$; we refer to [1] or to the introduction of [7]. Since the discovery of blown-up abelian surfaces in \mathcal{T} by Hirzebruch and Holzapfel some years ago (cf. [2]) there have been no further examples of surfaces of special type in \mathcal{T} . In this short note we present a new class of modular surfaces $X \in \mathcal{T}$ with $kod(X) = 1$.

In what follows we only consider complex projective smooth surfaces. Recall that a minimal elliptic surface $\pi : X \rightarrow C$ with finite Mordell-Weil group $MW(X)$ of sections is called *extremal* if $rank\ NS(X) = h^{1,1}(X)$. Particular examples arise in the following way. To each pair of positive integers

$$(m, n) \notin \{(1, 1), (1, 2), (2, 2), (1, 3), (1, 4), (2, 4)\}$$

there exists a modular elliptic surface over $\overline{\mathbb{Q}}$ in the sense of Shioda [10]

$$\pi_n(m) : X_n(m) \rightarrow C_n(m)$$

such that $\pi_n(m)$ admits no multiple fibers and has a non-constant j -invariant. By [10], $X_n(m)$ is an extremal elliptic surface with the following properties.

- $MW(X_n(m)) = \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$

2000 *Mathematics Subject Classification.* Primary 11J25; Secondary 14G35.

Key words and phrases. special surfaces, modular and Shimura varieties, Picard modular surfaces.

- $C_n(m)$ is the (compactified) curve $\overline{\Gamma_m(n) \backslash \mathbb{H}}$ where $\Gamma_n(m) \subset \mathbf{SL}_2(\mathbb{Z})$ is the group

$$\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix}; \begin{pmatrix} a & b \\ c & d \end{pmatrix} \equiv \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix} \bmod m, b \equiv 0 \bmod n \right\}.$$

- $C_n(m)$ parametrizes triples $((E, e_E), x, y)$ of elliptic curves E with neutral element $e_E \in E(\mathbb{C})$ and elements $x \in E[m], y \in E[n]$ such that $|\mathbb{Z}x + \mathbb{Z}y| = mn$.
- All singular fibers of $\pi_n(M)$ are of type I_k in Kodaira's notation; they lie over the cusps of $c \in C_n(m)$. A representant of c in $\mathbb{Q} \cup \{\infty\}$ is stabilized by a matrix $\gamma \in \Gamma$ which is a $\mathbf{SL}_2(\mathbb{Z})$ -conjugate of

$$\begin{pmatrix} 1 & k \\ 0 & 1 \end{pmatrix}.$$

The points x and y arise from the intersection of E with generators of $MW(X_n(m))$. More generally, by [8, Thm. 1.2, Thm. 1.3] each extremal elliptic surface $\pi : X \rightarrow C$ with non-constant j -invariant, no multiple fibers and $MW(\tilde{X}) = \mathbb{Z}/m\mathbb{Z} \times \mathbb{Z}/n\mathbb{Z}$, (m, n) as above, allows a cartesian diagram of isogenies

$$\begin{array}{ccc} X & \xrightarrow{\pi} & C \\ \downarrow & & \downarrow v \\ X_n(m) & \xrightarrow{\pi_n(m)} & C_n(m) \end{array}$$

With this perspective we are able to formulate our main result. We call a complex projective smooth surface X *irregular* if $h^1(X, \mathcal{O}_X) > 1$.

Theorem 1.1. *Let X be an irregular minimal surface in \mathcal{T} with $\text{kod}(X) = 1$ and finite Mordell-Weil group. Then after an étale base change X becomes an extremal elliptic surface fibered over an elliptic curve C such that the following assertions hold.*

- (1) *There is a cartesian diagram over $\overline{\mathbb{Q}}$*

$$\begin{array}{ccc} X & \xrightarrow{\pi} & C \\ \downarrow & & \downarrow \\ X_6(6) & \xrightarrow{\pi_6(6)} & C_6(6) \end{array}$$

- (2) *The compactification divisor D of X consists of 36 sections of π , each having self-intersection number $-\chi(X)$. The fibration π admits $2\chi(X)$ singular fibers of type I_6 , and each component of an I_6 intersects D in precisely 6 points. We have $\text{rank } NS(X) = 10\chi(X) + 2$.*

Conversely, $X_6(6)$ is an extremal elliptic and irregular surface in \mathcal{T} .

The diagram in the first statement is induced by the j -invariant.

2. SOME BASIC PROPERTIES OF SURFACES IN \mathcal{T}

We cite two results on ball-quotient surfaces which will be needed for the proof of the theorem. The first result is essentially [9, Thm. 3.1] specified to $\dim X = 2$ with attention to sign conventions, except the assertion on semi-stability. The latter assertion follows from [5]. A reduced effective divisor is called *semi-stable* if

it has normal crossings and if every rational smooth prime component intersects the remaining components in more than one point.

Theorem 2.1 (Tian-Yau/Miyaoka-Sakai). *Let X be a smooth projective surface and $D \subset X$ a divisor with normal crossings. Suppose that $K_X + D$ is big and ample modulo D . Then*

$$c_1^2(\Omega_X^1(\log D)) \leq 3c_2(\Omega_X^1(\log D)),$$

with equality holding if and only if $X \setminus D$ is an unramified ball quotient $\Gamma \setminus \mathbf{B}$ and D is semi-stable.

There is a canonical exact sequence

$$0 \longrightarrow \Omega_X^1 \longrightarrow \Omega_X^1(\log D) \xrightarrow{\text{res}} \mathcal{O}_D \longrightarrow 0$$

where res is the Poincaré residue map. With this one proves that $c_1(\Omega_X^1(\log D)) = [D] - c_1(X) \in H^2(X, \mathbb{C})$ and $c_2(\Omega_X^1(\log D)) = c_2(X) - (c_1(X), [D]) + ([D], [D]) \in H^4(X, \mathbb{C})$. Therefore,

$$c_1^2(\Omega_X^1(\log D)) = (K_X + D)^2.$$

It is interesting to note that if equality holds in the theorem, then D is smooth. Namely, if $\Gamma' \subset \Gamma$ is a neat normal subgroup with finite index in Γ , then $\Gamma' \setminus \mathbf{B}$ is compactified by a smooth elliptic divisor, and $\Gamma \setminus \mathbf{B}$ is compactified by a divisor D . As D is the quotient D'/G , $G = \Gamma/\Gamma'$, it is a normal curve. Hence, D is smooth and consists of elliptic curves, for rational curves cannot appear because of semi-stability. The next is proved *verbatim* as [7, Lemma 3.2].

Lemma 2.2. *Let X be in \mathcal{T} with compactification divisor D and consider an irreducible curve $L \subset X$. If L is smooth rational then $|L \cap D| \geq 3$. If L is a smooth elliptic curve then $|L \cap D| \geq 1$.*

3. PROOF OF THE RESULTS

General theory asserts that X admits an elliptic fibration $\pi : X \longrightarrow C$ which is the Albanese morphism. As $K_X + D$ is ample modulo D , it follows that a general fiber F has positive self-intersection with D . Thus, a component of D dominates C . Hence, C is an elliptic curve and $h^1(X, \mathcal{O}_X) = 1$. Moreover, after transition to an étale cover \tilde{C} of C and performing a base change, we can achieve that every D_i is a section, as soon as it dominates C ([7, Lemma 3.3]). We will assume this from now on. Since the curves D_i are pair-wise disjoint, they must be all sections.

Lemma 3.1. *The identities $36\chi(X) = DF \cdot \chi(X) = -D^2$ and $DF = 36$ hold.*

Proof. The canonical bundle-formula implies that $K_X = \pi^*(\mathfrak{c})$ with a divisor Weil divisor $\mathfrak{c} \in \text{Div}(C)$. Moreover, $h^0(X, mK_X) = h^0(C, m\mathfrak{c})$. The theorem of Riemann-Roch yields $h^0(X, K_X) = \deg \mathfrak{c} > 0$. Adjunction formula implies that

$$D_i^2 = -\deg \mathfrak{c} = -h^0(X, K_X) = -\chi(X).$$

Hence, $-D^2 = -\sum D_i^2 = DF\chi(X)$. Furthermore, $12\chi(X) = c_2(X)$ by Noether's formula. So, Thm. 2.1 yields the remaining identities. \square

We consider the Mordell-Weil group $MW(X) = MW_{\text{tor}}(X)$. It follows that $|MW_{\text{tor}}(X)| \geq 36$. We prove the following lemma of general interest.

Lemma 3.2. *Let $\pi : X \rightarrow C$ be a minimal elliptic surface over an elliptic curve C and assume that $\text{kod}(X) \geq 1$ and that each rational curve $L \subset X$ meets at least three sections of π . Suppose moreover that $D = MW_{\text{tor}}(X) \geq 33$. Then all singular fibers of π are semi-stable of type I_6 , X has $2\chi(X)$ singular fibres and $MW(X) = MW_{\text{tor}}(X) = \mathbb{Z}/6\mathbb{Z} \times \mathbb{Z}/6\mathbb{Z}$ and the rank of the Neron-Severi group $NS(X)$ equals $h^{1,1}(X) = 10\chi(X) + 2$.*

Proof. The assertion concerning $MW(X)$ follows directly from [4, (4.8)]. [4, Lemma 1.1] implies then that all singular fibers are of type I_n . If $H_n \subset M(X)$ is the non-trivial isotropy group of a node $x \in I_n$ then $MW_{\text{tor}}(X)/H_n$ is cyclic by [4, Lemma 2.2]. Moreover, all nodes from one and the same fiber have the same isotropy group by [4, Lemma 2.1, (c)], and this isotropy group is non-trivial by [4, Lemma 2.1, (b)] and because a component of I_n meets at least three sections. Thus, always $|H_n| \geq 6$. On the other hand, by [4, p. 251] and [4, Lemma 2.3, (f)], $\sum_{I_n} n = c_2(X)$ and

$$36c_2(X) = |MW_{\text{tor}}(X)|c_2(X) = \sum_{I_n} n|H_n|^2.$$

Hence, always $|H_n| = 6$. Let $S \in MW(X)$ be the neutral element. By the proof of [4, Lemma 2.2], H_n consists of precisely those sections meeting the prime component $L \subset I_n$ which contains $S \cap I_n$. However, since we may take any section to be the neutral element of $MW(X)$, for each component $L \subset I_n$ we have $LD = 6$. As $DI_n = 36$, we get $n = 6$. Finally, recalling that $\sum_{I_n} n = c_2(X)$, we find for the number t of singular fibers:

$$t = 2\chi(X) = 2g(C) - 2 + \text{rank } MW(X) + 2\chi(X).$$

According to [4, Prop. 1.6] this equality holds if and only if $\text{rank } NS(X) = h^{1,1}(X)$. An easy calculation shows now that $h^{1,1}(X) = 10\chi(X) + 2$. \square

It follows that X is isomorphic to a pull-back $X_6(6) \times_{C_6(6)} C$.

REFERENCES

- [1] Holzapfel, R.-P., *Ball and Surface Arithmetics*, Aspects vol. **E29** (Vieweg, Braunschweig, 1998)
- [2] Holzapfel, R.-P., *Jacobi theta embedding of a hyperbolic 4-space with cusps*, in: Geometry, integrability and quantization (Coral Press Sci. Publ., Sofia, 2002).
- [3] Kodaira, K., *On stability of compact submanifolds of compact complex manifolds*, Am. J. Math. vol. **85** (1963), 79-94
- [4] Miranda, R.; Persson, U., *Torsion groups of elliptic surfaces*, Comp. Math. tome **72** (1989), 249-267
- [5] Miyokawa, Y., *The maximal number of quotient singularities on surfaces with given numerical invariants*, Math Ann. **268** (1984)
- [6] Mostow, G.D., *Strong Rigidity of Locally Symmetric Spaces*, Annals of Mathematics Studies, No. **78** (Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1973)
- [7] Momot, A., *Irregular ball-quotient surfaces with non-positive Kodaira dimension*, to appear
- [8] Kloostermann, R., *Extremal elliptic surfaces and infinitesimal Torelli*, Michigan Math. J. 52 (2004)
- [9] Tian, G., Yau, S.T., *Existence of Kähler-Einstein metrics on complete Kähler manifolds and their applications to algebraic geometry*, in: Yau (ed.), Mathematical Aspects of String Theory, Advanced Series in Mathematical Physics vol.1 (World Scientific, Singapore, 1987)
- [10] Shioda, T., *On elliptic modular surfaces*, J. Math. Soc. Japan 24 (1972), 20-59

DEPARTEMENT MATHEMATIK, ETH ZÜRICH, HG J65, RÄMISTRASSE 101, 8092 ZÜRICH, SWITZER-
LAND

E-mail address: `aleksander.momot@math.ethz.ch`